

Altered Mental Status

Note:
Request Advanced Life Support if available.
Do Not delay transport to the appropriate hospital.

Note:
This protocol is for patients who are not alert (A), but who are responsive to verbal stimuli (V), responding to painful stimuli (P), or unresponsive (U).

- I. Assess the situation for potential or actual danger. If the scene/situation is not safe, retreat to a safe location, create a safe zone and obtain additional assistance from a police agency.

Note:
Emotionally disturbed patients must be presumed to have an underlying medical or traumatic condition causing the altered mental status.

Note:
All suicidal or violent threats or gestures must be taken seriously. These patients should be in police custody if they pose a danger to themselves or others.
If the patient poses a danger to themselves and/or others, summon police for assistance.

- II. Perform primary assessment. Assure that the patient's airway is open and that breathing and circulation are adequate. Suction as necessary
- III. Administer high concentration oxygen. In children, humidified oxygen is preferred.
- IV. Obtain and record patient's vital signs, including determining the patient's level of consciousness. Assess and monitor the Glasgow Coma Scale.
 - A. **If the patient is unresponsive (U) or responds only to painful stimuli (P), prepare for transport** while continuing care.

Altered Mental Status, continued

B. If the patient has a known history of diabetes controlled by medication, is conscious and is able drink without assistance, provide an oral glucose solution, fruit juice or non-diet soda by mouth, then transport, keeping the patient warm. If regionally approved to obtain blood glucose levels utilizing a glucometer, follow your regionally approved protocol.

C. If patient has a suspected narcotic overdose:

- i.** Respirations less than 10/minute and signs of respiratory failure or respiratory arrest, refer to appropriate respiratory protocol.
- ii.** If regionally approved and available, obtain patient's blood glucose (BG) level.
 - 1.** If BG is less than 65, follow IV.B above.
 - 2.** If BG is more than 65, proceed to next step.
- iii.** Administer 2mg/2ml of naloxone (Narcan®) via a mucosal atomizer device (MAD).

1. Exclusion criteria:

- a.** Cardiopulmonary Arrest,
 - b.** Seizure activity during this incident,
 - c.** Pediatric patients,
 - d.** Therapeutic opiate use through physician prescription,
 - e.** Evidence of nasal trauma, nasal obstruction and/or epistaxis.
- 2.** Insert MAD into patient's left nostril and inject 1mg/1ml.
 - 3.** Insert MAD into patient's right nostril and inject 1mg/1ml.
 - 4.** Prepare for transport. After 5 minutes, if patient's respiratory rate is not greater than 10 breaths/minute, administer a second dose of naloxone 2mg/2ml follow the same procedure as above.

Note:

Do not give solutions by mouth to patients who are unconscious or to patients with head injuries.

- V.** If underlying medical or traumatic condition causing an altered mental status is not apparent; the patient is fully conscious, alert (A) and able to communicate; and an emotional disturbance is suspected, proceed to the Behavioral Emergencies protocol.

Altered Mental Status, continued

- VI. Transport to the closest appropriate facility while re-evaluating vital signs every 5 minutes and reassess as necessary.
- VII. Record all patient care information, including the patient's medical history and all treatment provided, on a Prehospital Care Report (PCR).